

(Cardiovascular and Cerebrovascular) Analysis Report Card

Name: Example(Female)

Sex: Female

Age: 30

Figure: Standard body weight(165cm,62kg)

Testing Time: 2013-05-02 17:20

Actual Testing Results

Testing Item	Normal Range	Actual Measurement Value	Testing Result
Blood Viscosity	48.264 - 65.371	66,966	
Cholesterol Crystal	56.749 - 67.522	63,236	
Blood Fat	0.481 - 1.043	1,237	
Vascular Resistance	0.327 - 0.937	1,709	
Vascular Elasticity	1.672 - 1.978	1,431	
Myocardial Blood Demand	0.192 - 0.412	0,503	
Myocardial Blood Perfusion Volume	4.832 - 5.147	4,646	
Myocardial Oxygen Consumption	3.321 - 4.244	5,485	
Stroke Volume	1.338 - 1.672	0,927	
Left Ventricular Ejection Impedance	0.669 - 1.544	1,505	
Left Ventricular Effective Pump Power	1.554 - 1.988	0,872	
Coronary Artery Elasticity	1.553 - 2.187	2,095	
Coronary Perfusion Pressure	11.719 - 18.418	12,296	
Cerebral Blood Vessel Elasticity	0.708 - 1.942	1,66	
Brain Tissue Blood Supply Status	6.138 - 21.396	20,5	

Reference Standard:

Normal(-)	Mildly Abnormal(+)
Moderately Abnormal (++)	Severely Abnormal (+++)

Blood Viscosity:	48.264-65.371(-)	65.371-69.645(+)
	69.645-73.673(++)	>73.673(+++)
Cholesterol Crystal:	56.749-67.522(-)	67.522-69.447(+)

	69.447-74.927(++)	>74.927 (+++)
Blood Fat:	0.481-1.043(-) 1.669-1.892(++)	1.043-1.669(+) >1.892(+++)
Vascular Resistance:	0.327-0.937(-) 1.543-1.857(++)	0.937-1.543(+) >1.857(+++)
Vascular Elasticity:	1.672-1.978(-) 1.511-1.047(++)	1.672-1.511(+) <1.047(+++)
Myocardial Blood Demand:	0.192-0.412(-) 0.571-0.716(++)	0.412-0.571(+) >0.716(+++)
Myocardial Blood Perfusion Volume:	4.832-5.147(-) 4.029-4.177(++)	4.177-4.832(+) <4.029(+++)
Myocardial Oxygen Consumption:	3.321-4.244(-) 5.847-6.472(++)	4.244-5.847(+) >6.472(+++)
Stroke Volume:	1.338-1.672(-) 0.139-0.647(++)	0.647-1.338(+) <0.139(+++)
Left Ventricular Ejection Impedance:	0.669-1.544(-) 2.037-2.417(++)	1.544-2.037(+) >2.417(+++)
Left Ventricular Effective Pump Power:	1.554-1.988(-) 0.597-1.076(++)	1.076-1.554(+) <0.597(+++)
Coronary Artery Elasticity:	1.553-2.187(-) 0.983-1.182(++)	1.182-1.553(+) <0.983(+++)
Coronary Perfusion Pressure:	<8.481(+++) 18.418-21.274(++)	8.481-11.719(++) >21.274(+++)
Cerebral Blood Vessel Elasticity:	0.708-1.942(-) 0.109-0.431(++)	0.431-0.708(+) <0.109(+++)
Brain Tissue Blood Supply Status:	6.138-21.396(-) 1.214-3.219(++)	3.219-6.138(+) <1.214(+++)

Parameter Description

Blood Viscosity(N): The basic indicator of Hemorheology refers to the internal friction among blood molecules.

Hyperviscosity state: Namely, the viscosity of blood is high, and blood flow is affected. Therefore, high blood pressure patients with high viscosity are prone to have cerebrovascular accidents, such as stroke and other phenomena; coronary heart disease patients with high viscosity are prone to have myocardial infarction and so on.

The blood flow in the blood vessels is in a laminar flow state, which is stratified flow. The flow

velocity close to the vessel wall is slower, and the flow velocity is fastest in the middle. Thus, the larger the shear rate is, the greater the slope is, the greater the shear stress is, the faster the flow velocity is, and the lower the N is. The smaller the shear rate is, the lower the slope is, the smaller the shear stress is, the lower the flow velocity is, and the higher the N is.

Cholesterol Crystal:

(1) Increase is seen in primary high cholesterol blood, the aura of mild atherosclerosis, blood stagnation type chest pain, phlegm congestion type chest pain, etc.

(2) Reduction is seen in decreased immunity, malnutrition, cardiac insufficiency, Qi and Yin deficiency type chest pain, Yang Qi deficiency type chest pain, etc.

Blood Fat:Blood fat abnormality is divided into primary abnormality and secondary abnormality.

1. Primary Hyperlipoproteinemia: refers to hyperlipoproteinemia caused by the possibility of unknown cause related to certain environmental factors (including diet, nutrition, drugs, etc.), or gene mutations.

2. Secondary Hyperlipoproteinemia: refers to hyperlipidemia caused by certain systemic diseases or drugs, such as hyperlipidemia caused by diabetes, hypothyroidism, nephrotic syndrome, chronic renal failure and acute renal failure and so on.

(1) Increase is seen in idiopathic hyperlipidemia, atherosclerosis, blood stagnation type chest pain, etc.

(2) Reduction is seen in ferrite decreased immunity, the Qi and Yin deficiency type chest pain, etc.

(3) Decline is seen in decline of cerebral arterial oxygen content and mild ischemic cerebrovascular disease aura.

Vascular Resistance:

Increase is in direct proportion to the length of blood vessels, and is in inverse proportion to the caliber of blood vessels. The increase of vascular resistance is seen in mildly elevated systolic and diastolic blood pressure, mild hypertension, insomnia with deficiency of both heart and spleen, phlegm-heat internal confusion type insomnia, etc.

Decline is seen in mildly declined systolic and diastolic blood pressure, mild hypotension, Yin deficiency and Huo exuberance type insomnia, etc.

Vascular Elasticity:refers to the expansion extent of arterial vascular elasticity during systolic ejection.

Influence Factors: (1) The size of SV. The greater the SV is, the greater the FEK is. (2) Emptying rate. The faster the emptying rate is, the smaller the FEK is. (3) Bad vascular elasticity.

The SV is not low, the emptying rate is not fast, and the FEK is also small, so it is possible to determine the possibility of hardening of blood vessels. It should not determine the possibility by a single parameter. The increase of vascular elasticity is seen in the mildly elevated systolic blood pressure, the mildly reduced diastolic blood pressure, the mildly increased pulse press and slightly higher blood pressure. The decline is seen in mildly atherosclerosis, coronary heart diseases, blood stagnation type chest pain, Yang Qi deficiency type chest pain, etc.

Myocardial Blood Demand:The blood demand per minute of coronary artery perfusion of heart.

Myocardial Blood Perfusion Volume:The actual blood demand per minute of coronary artery perfusion of heart.

Myocardial Oxygen Consumption:The milliliter value of oxygen consumption of heart per minute.

Influence Factors: Three aspects

(1) Heart rate: the heart rate is fast, and the HOV is great.

(2)(2) Myocardial contractility: the cardiac contractility is strong, and the HOV is great.

(3) Myocardial contraction time: the longer the contraction time is, the greater the HOV is.

Thus, low oxygen consumption and high cardiac work are the best state.

Stroke Volume:The blood volume output by the heart in beat each time.

Influence Factors: Five aspects

(1) The effective circulating blood volume (BV): when the blood volume is insufficient, the returned blood volume is little, and the SV is reduced.

(2) The weakening of myocardial contractility: the contractility is low, and the pressure is low, so the ejected blood volume is less.

(3) The extent of ventricular filling: In range of myocardial elasticity, the greater the degree of filling is, the stronger the retraction is, and the SV is increased. The normal heart chamber capacity is 173ml, but not all of the blood is ejected. The blood volume in the left ventricle is about 60% -70% of the total capacity, being about 125ml or so. Usually, the Chinese people's average SV is 80-90ml.

(4) The size of peripheral vascular resistance (PR). The PR is large, and then the SV is reduced; the PR is small, and then the SV is increased.

(5) Ventricle wall movement.

When the ventricle is contracted, the cardiac muscle is in coordinated movement. If the myocardial contraction is not coordinated, the SV is reduced. For instance, some patients with myocardial infarction have part of infarction, so the myocardial contractility is inconsistent and the SV is reduced. However, under normal circumstances, the ventricle wall movement can not be abnormal.

Left Ventricular Ejection Impedance:reflects the indicators of resistance status of the left ventricular outflow channel.

Influence Factors:

(1) The fact whether the outflow channel has lesion. The aortic stenosis and other conditions can make VER increased.

(2) The outflow channel has no lesion, while the emptying rate of aortic blood is slow, so VER is increased.

(3) The entire vascular resistance is large.

Left Ventricular Effective Pump Power:reflects the contraction strength of effective stroke of blood of the left ventricle.

Normally, the people: 1.8 kilograms. Pump power is low, and contraction is not good, so myocardial fibers may have problems. Pump power is high, and contractility is good, so the ejected blood volume is much.

Influence Factors: Four aspects

(1) The extent of ventricular filling: In range of elasticity, the greater the degree of filling is, the stronger the contractility is; the degree of filling and the contractility are in direct proportion. If out of the limit, the myocardial expansion is large, but the contractility is reduced. Thus, the proper degree of filling is a factor influencing the contractility.

(2) The effective circulating blood volume (returned blood volume BV): The returned blood volume is little, the filling is insufficiency, and the contractility is small; the returned blood volume is much, the filling is better, and the contractility is strong.

(3) The functional status of myocardium itself: The fact whether the myocardium has lesion. For instance, myocarditis. Myocardial cells are damaged, and myocardial elasticity is reduced, so the contractility is lowered.

(4) The normal degree of blood and oxygen supply of myocardium itself: The blood and oxygen supply is insufficient, so the contractility is lowered. Myocardial Oxygen Consumption: the milliliter value of oxygen consumption of heart per minute.

Coronary Artery Elasticity:

The source of power of life is the heart, and the blood nourishing the body constantly flows under her impetus. However, she also demands the nourishing of blood. Coronary artery, namely three blood vessels respectively located in the heart, can supply blood and oxygen to her. The coronary artery is the artery special for supplying blood to the heart. If cholesterol and other substances are accumulated in the blood vessels, the vascular cavity will be narrower or be blocked and the blood flow will be smooth and then be blocked to cause cardiac ischemia and a series of symptoms which are coronary heart disease, namely coronary atherosclerosis. Coronary heart disease is also called as coronary atherosclerotic heart disease. The excessive fat deposition results in atherosclerosis and weakened elasticity. The mortality of human on cardiovascular and cerebrovascular diseases induced on the arterial vessel wall has exceeded 1 / 2 of the total mortality of population.

Dangerous factors making the elasticity of coronary artery weakened: high blood fat, smoking, diabetes, obesity, high blood pressure, lack of physical activity, Psychological overstrain, family

history of coronary heart disease, oral contraceptive, etc.

Coronary Perfusion Pressure:the pressure of coronary artery of heart in blood supply is influenced by diastolic blood pressure and left atria pressure.

Part of myocardial ischemia, insufficient myocardial blood supply and entire myocardial ischemia can lead to myocardial infarction.

Cerebral Blood Vessel Elasticity:

The brain artery or the neck artery controlling the brain has lesion, which leads to disorder of intracranial blood circulation and damage of brain tissue. The elasticity of hardened brain blood vessels is weakened, and the vessel cavity is narrowed, so it is easy to form cerebral thrombosis. After the patients with cerebral arteriosclerosis excessively drink, the blood pressure will be suddenly elevated, the blood vessels will ruptured, so it is prone to form cerebral hemorrhage. After load drinking of alcohol, the concentration of alcohol in blood can reach its peak in a half hour. The alcohol can not only directly stimulate the blood vessel wall to make it lose its elasticity but also stimulate the liver to promote the synthesis of cholesterol and triglyceride,thus leading to atherosclerosis and cerebral atherosclerosis. Cerebrovascular disease can be divided into acute cerebrovascular disease and chronic cerebrovascular disease according to their process. The acute cerebrovascular disease includes trans ient ischemic attack, cerebral thrombosis,cerebral embolism, hypertensive encephalopathy, cerebral hemorrhage, subarachnoid hemorrhage, etc. The chronic cerebrovascular disease includes cerebral arteriosclerosis, cerebrovascular dementia, cerebral artery steal syndrome, Parkinson's disease, etc. The cerebrovascular disease which is known generally refers to the acute cerebrovascular disease. It often endangers the human life due to acute incidence, so it is easy to cause the attention. The chronic cerebrovascular disease is easy to be ignored by people due to its long course.

Brain Tissue Blood Supply Status:

Brain tissue blood supply mainly depends on the brain artery or the neck artery controlling the brain. Cerebrovascular diseases can be divided into two categories according to their nature, one is the ischemic cerebrovascular disease and the other one is the hemorrhagic cerebrovascular disease. There are many cases about the ischemic cerebrovascular disease in clinic, the patients account for 70% ~ 80% of all patients with cerebrovascular disease. Due to cerebral arteriosclerosis and other reasons, the vessel cavity of brain artery is narrowed, the blood flow is reduced or completely blocked, the brain blood circulation is disordered, and the brain tissue is damaged,so a series of symptoms occur. The hemorrhagic cerebrovascular disease is mainly caused by long-term high blood pressure, congenital cerebral vascular malformation and other factors.Due to blood vessel rupturing, blood spilling, oppression on brain tissue and blocked blood circulation, the patients often show increased intracranial pressure, disorientation and other symptoms. Thus, the patients account for about 20% ~ 30% of all patients with cerebrovascular disease.

The test results for reference only and not as a diagnostic conclusion.